

Reg.No										
--------	--	--	--	--	--	--	--	--	--	--



MANIPAL UNIVERSITY, MANIPAL

FIRST SEMESTER M.SC (APPLIED MATHEMATICS & COMPUTING) MAKE UP EXAMINATION,

**SUB: DIFFERENTIAL EQUATIONS(MAT - 601)
(REVISED CREDIT SYSTEM)**

Time : 3 Hrs.

Max.Marks : 50

Note : a) Answer any FIVE full questions. b) All questions carry equal marks (3+ 3+ 4).

- 1A. If $y = x$ is one of the solution of the differential equation $(x^2 + 1) y'' - 2 x y' + 2 y = 0$. Find the general solution of $(x^2 + 1) y'' - 2 x y' + 2 y = 6(x^2 + 1)^2$.
- 1B. Obtain the Rodrigue's formula $P_n(x) = \frac{1}{n! 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n$ where $P_n(x)$ is Legendre polynomial of degree n .
- 1C. State and prove uniqueness theorem for Let $L(y) = y^{(n)} + a_1 y^{(n-1)} + a_2 y^{(n-2)} + \dots + a_n y = 0$ on an interval I containing a point x_0 . With usual initial conditions
- 2A. Find a function $\varphi(x)$ which has a continuous derivative on $0 \leq x \leq 2$ which satisfies $\varphi(0) = 0$, $\varphi'(0) = 1$ and $y'' - y = 0$ for $0 \leq x \leq 1$ and $y'' - 9y = 0$ for $1 \leq x \leq 2$.
- 2B. Prove that (i) $J_{5/2}(x) = \sqrt{\frac{2}{\pi x}} \left\{ \frac{3-x^2}{x^2} \sin x - \frac{3}{x} \cos x \right\}$
(ii) $J_n''(x) = \frac{1}{4} [J_{n-2}(x) - 2J_n(x) + J_{n+2}(x)]$
- 2C. Solve by annihilator method solve $y'' + y' = 4x^2 e^x$
- 3A Test the solution functions φ_1, φ_2 , defined by $\varphi_1 = x^2$, $\varphi_2 = x|x|$, for linearly independent on $-\infty < x < \infty$.
- 3B. Solve in series $x \frac{d^2 y}{dx^2} + \frac{dy}{dx} + xy = 0$.

- 3C Prove that (1) $H_n'(x) = 2n H_{n-1}(x)$
 (2) $2x H_n(x) = 2n H_{n-1}(x) + H_{n+1}(x)$
- 4A. If $\phi_1(x)$ is a solution of a differential equation $y'' + a_1(x) y' + a_2(x) y = 0$ then show that $\phi_2(x) = \phi_1(x) f(x)$ is a solution of this equation provided $f'(x)$ satisfies the equation $(\phi_1^2 y)' + a_1(x) (\phi_1^2 y) = 0$
- 4B Find the eigen values and the eigen functions of $y'' + \lambda y = 0$, $y(0) = y(\pi) = 0$.
- 4C If ϕ_1, ϕ_2 are two solutions of $L(y) = 0$ on an interval I containing a point x_0 then prove that $W(\phi_1, \phi_2)(x) = e^{-\int a_1(x-x_0) dx} W(\phi_1, \phi_2)(x_0)$
- 5A. Solve by method of variation of parameters $y''' + y'' + y' + y = 1$
- 5B.(i) Show that $\int_{-1}^1 x^2 P_{n-1}(x) P_{n+1}(x) dx = \frac{2n(n+1)}{(2n-1)(2n+1)(2n+3)}$
- 5C. Find the solution of $x^2 y'' + 9x y' + 12y = 0$ by series method.
- 6A. Obtain the series solution of $(1-x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + n(n+1)y = 0$, n is a real number.
- 6B. Let ϕ_1, ϕ_2 be two solutions of $L(y) = y'' + a_1 y' + a_2 y = 0$. Then prove that ϕ_1, ϕ_2 are linearly independent on an interval I if and only if $W(\phi_1, \phi_2)(x) \neq 0$.
- 6C. Prove that $\int_0^1 x J_n(\alpha x) J_n(\beta x) dx = \begin{cases} 0 & , \alpha \neq \beta \\ \frac{1}{2} [J_{n+1}(\alpha)]^2 & , \alpha = \beta \end{cases}$ where α, β are the roots of $J_n(x) = 0$.