



MANIPAL
ACADEMY of HIGHER EDUCATION

(Deemed to be University under Section 3 of the UGC Act, 1956)

Reg. No.

--	--	--	--	--	--	--	--	--	--

**DEPARTMENT OF SCIENCES, M. Sc. PHYSICS
I SEMESTER END SEMESTER EXAMINATIONS
NOVEMBER 2018**

**Subject: Quantum Mechanics I (PHY-4105)
(REVISED CREDIT SYSTEM)**

Time: 3 Hours Date: November 2018 MAX. MARKS: 50

**Note: (i) Answer all the questions.
(ii) Answer the questions to the point.**

1. (i) An electron has a de Broglie wavelength of 1.5×10^{-12} m. Find its (a) kinetic energy and (b) group and phase velocities of its matter waves. [5]

(ii) Show that the average kinetic energy of a particle of mass m with a wave function $\psi(x)$ can be written in the form

$$T = \frac{\hbar^2}{2m} \int_{-\infty}^{\infty} \left| \frac{d\psi}{dx} \right|^2 dx = 0 \quad [5]$$

2. (i) Write the time dependent Schroedinger equation. Using separation of variable technique split it into purely time dependent and time independent components. [3]

(ii) What are stationary states? [2]

(iii) Show that the probability density of the linear harmonic oscillator in an arbitrary superposition state is periodic with the period equal to the period of the oscillator. [5]

3. (i) Calculate the energy eigenvalues and eigenfunctions of a particle confined in a two dimensional square shaped infinite deep potential well. Show that the eigenstates are degenerate. [5]

(ii) Calculate the energy difference between the stationary states $l = 1$ and $l = 2$ of the rigid molecule H_2 . Use the Bohr frequency rule to estimate the frequency of radiation involved during transition between these two states. Suggest a method for determining the bond length of hydrogen molecule. [5]

4. Solve the Schroedinger equation for the hydrogen atom and discuss the radial wavefunction. [10]

5. (i) Prove that it is impossible to construct a completely antisymmetric spin function for three electrons. [5]

(ii) Write a short note on exchange operators. Calculate their eigenvalues. [5]

Useful formulae:

$$\nabla^2 t = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial t}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial t}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 t}{\partial \phi^2}$$

$$\int_0^\infty \exp(-a^2 x^2) \cos(bx) dx = \frac{\sqrt{\pi}}{2a} \exp\left(-\frac{b^2}{4a^2}\right)$$

$$\int_0^\infty x^n \exp(-ax) dx = \frac{n!}{a^{n+1}}, \quad \text{where } n \geq 0, \quad a > 0$$